



令和8年度 フェロモントラップ等調査結果 セジロウンカ(予察灯)

1 調査結果

(1) グラフ



(2)調査データ



地帯区分		中西部			南部		
設置場所		東広島市八本松原			呉市安浦町		
周辺作物		水稻			水稻		
月	半旬	本年	平年	前年	本年	平年	前年
5月	1	0.0	0	0	0.0	0	0
	2	0.0	0	0	0.0	0	0
	3	0.0	0	0	0.0	0	0
	4	0.0	0	0	0.0	0	0
	5	0.0	0	0	0.0	0	0
	6	0.0	0	0	0.0	0.1	0
6月	1	0.0	0	0	0.0	0	0
	2	0.0	0	0	0.0	0.1	1
	3	0.0	0.1	0	0.0	0.1	1
	4	0.0	0	0	0.0	1.6	0
	5	0.0	0.4	4	0.0	6.2	1
	6	0.0	1.8	0	1.0	4.0	0
7月	1	21.0	2.3	0	31.0	3.0	0
	2	27	0.6	0	11.0	1.1	0
	3	0	2.3	1	0	5.8	0
	4	0	1.1	0	1.0	3.9	0
	5	0	1.4	0	1.0	1.6	0
	6	1	0.6	0	1.0	1.2	0
8月	1	0	0.9	0	1.0	0.8	2
	2	0	2.1	7	0	2.3	12
	3	0	1.3	1	0	7.7	7
	4	0	2.7	1	1.0	9.6	1
	5	0	1.3	0	0	2.0	3
	6	0	1.4	0	0	13.5	10
9月	1	0	7.9	0	0	4.5	1
	2		2.6	0	0	1.9	2
	3		3.0	0	2.0	2.7	1
	4		0.6	0		6.0	1
	5		0.9	0		1.3	0
	6		1.1	4		2.0	0

安浦:5/27～29欠測

令和8年度 フェロモントラップ等調査結果 トビイロウンカ(予察灯)

1 調査結果

(1) グラフ



(2)調査データ

地帯区分		中西部			南部		
設置場所		東広島市八本松原			呉市安浦町		
周辺作物		水稻			水稻		
月	半旬	本年	平年	前年	本年	平年	前年
5月	1	0	0	0	0	0	0
	2	0	0	0	0	0	0
	3	0	0	0	0	0	0
	4	0	0	0	0	0	0
	5	0	0	0	0	0	0
	6	0	0	0	0	0	0
6月	1	0	0	0	0	0	0
	2	0	0	0	0	0	0
	3	0	0	0	0	0	0
	4	0	0	0	0	0	0
	5	0	0	0	0	0.1	0
	6	0	0	0	0	0	0
7月	1	0	0	0	0	0	0
	2	2.0	0	0	0	1.3	1.0
	3	0	0	0	0	2.2	0
	4	0	0	0	0	0.2	0
	5	0	0	0	0	0.2	0
	6	0	1	0	0	0	0
8月	1	0	0	0	0	0	0
	2	0	0	0	0	0.5	0
	3	0	0	0	0	0.6	1.0
	4	0	1	0	0	0.4	0
	5	4.0	0	0	1.0	0.4	0
	6	0	1.1	0	0	10.1	0
9月	1	0	19.9	0	0	46.8	1.0
	2		8.5	0	0	22.9	0
	3		4.4	0	3.0	14.9	0
	4		2.3	0		2.8	0
	5		0.5	0		0.8	0
	6		2.6	0		8.0	0

安浦:5/27～29欠測